
200600036

No.



THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

Holden's Joundation Seeds L. C.

DECEMP, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT'S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY FEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC PREMISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE HT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR TING IT, OR EXPORTING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE URPOSES, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT BY THE PLANT VARIETY PROTECTION ACT. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

CORN, FIELD

'LH391'

In Jestimony Thereof, I have hereunto set my hand and caused the seal of the Hant Baristy Frotestion Office to be affixed at the City of Washington, D.C. this twenty-ninth day of April, in the year two thousand and eight.

Allet

Denz-

Commissioner

Plant Variety Protection Offico Agricultural Marketing Servico dward T. Schafer

f Agriculturo

CAPACITY OR TITLE

DATE

(See reverse for instructions and information collection burden statemen

Timothy R. Kain

Patent Scientist

ST-470 (02-10-2003) designed by the Plant Variety Protection Office using Word 2000

CAPACITY OR TITLE

INSTRUCTIONS

GENERAL: To be effectively filed with the Plant Variety Protection Office (PVPO), ALL of the following items must be received in the PVPO: (1) Completed application form signed by the owner; (2) completed exhibits A, B, C, E; (3) for a seed reproduced variety at least 2,500 viable untreated seeds, for a hybrid variety at least 2,500 untreated seeds of each line necessary to reproduce the variety, or for tuber reproduced varieties verification that a viable (in the sense that it will reproduce an entire plant) tissue culture will be deposited and maintained in an approved public repository; (4) check drawn on a U.S. bank for \$3,652 (\$432) filing fee and \$3,220 examination fee), payable to "Treasurer of the United States" (See Section 97.6 of the Regulations and Rules of Practice.) Partial applications will be held in the PVPO for not more than 90 days, then returned to the applicant as unfiled. Mail application and other requirements to Plant Variety Protection Office, AMS, USDA, Room 401, NAL Building, 10301 Baltimore Avenue, Beltsville, MD 20705-2351. Retain one copy for your files. All items on the face of the application are self explanatory unless noted below. Corrections on the application form and exhibits must be initialed and dated. DO NOT use masking materials to make corrections. If a certificate is allowed, you will be requested to send a check payable to "Treasurer of the United States" in the amount of \$432 for issuance of the certificate. Certificates will be issued to owner, not licensee or agent.

Plant Variety Protection Office Telephone: (301) 504-5518 FAX: (301) 504-5291

Homepage: http://www.ams.usda.gov/science/pvpo/pvp.htm

ITEM

18a. Give:

- (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method;
- the details of subsequent stages of selection and multiplication;
- (3) evidence of uniformity and stability; and
- (4) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified
- 18b. Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all other varieties in the same crop. If the new variety is most similar to one variety or a group of related varieties:
 - (1) identify these varieties and state all differences objectively;
 - attach statistical data for characters expressed numerically and demonstrate that these are clear differences; and
 - (3) submit, if helpful, seed and plant specimens or photographs (prints) of seed and plant comparisons which clearly indicate distinctness.
- 18c. Exhibit C forms are available from the PVPO Office for most crops; specify crop kind. Fill in Exhibit C (Objective Description of Variety) form as completely as possible to describe your variety.
- 18d. Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant color, disease resistance, etc.
- 18e. Section 52(5) of the Act requires applicants to furnish a statement of the basis of the applicant's ownership. An Exhibit E form is available from the PVPO.
- 19. If "Yes" is specified (seed of this variety be sold by variety name only, as a class of certified seed), the applicant MAY NOT reverse this affirmative decision after the variety has been sold and so labeled, the decision published, or the certificate issued. However, if "No" has been specified, the applicant may change the choice. (See Regulations and Rules of Practice, Section 97.103).
- 22. See Sections 41, 42, and 43 of the Act and Section 97.5 of the regulations for eligibility requirements.
- See Section 55 of the Act for instructions on claiming the benefit of an earlier filing date.
- 21. CONTINUED FROM FRONT (Please provide a statement as to the limitation and sequence of generations that may be certified.)
- 22. CONTINUED FROM FRONT (Please provide the date of first sale, disposition, transfer, or use for each country and the circumstances, if the variety (including any harvested material) or a hybrid produced from this variety has been sold, disposed of, transferred, or used in the U.S. or other countries.)

Sold in U.S. - December 2004

23. CONTINUED FROM FRONT (Please give the country, date of filing or issuance, and assigned reference number, if the variety or any component of the variety is protected by intellectual property right (Plant Breeder's Right or Patent).)

NOTES: It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownership or assignment or owner's representative during the life of the application/certificate. There is no charge for filling a change of address. The fee for filling a change of ownership or assignment or any modification of owner's name is specified in Section 97.175 of the regulations. (See Section 101 of the Act, and Sections 97.130, 97.131, 97.175(h) of the Regulations and Rules of Practice.)

To avoid conflict with other variety names in use, the applicant must check the appropriate recognized authority. For example, for agricultural and vegetable crops, contact: Seed Branch, AMS, USDA, Room 213, Building 306, Beltsville Agricultural Research Center--East, Beltsville, MD 20705, Telephone: (301) 504-8089. http://www.ams.usda.gov/lsg/seed.htm

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 9581-9055. The time required to complete this information collection is estimated to average 3.0 hours per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

ST-470 (02-10-2003) designed by the Plant Variety Protection Office with Word 2000. Replaces former versions of ST-470, which are obsolete.

EXHIBIT A

Origin and Breeding History LH391

LH391 was developed from the cross of LH277 x LH185 by selfing and using the conventional ear-to-row system of plant breeding. Yield, stalk quality, root quality, disease tolerance, late plant greenness, late plant intactness, ear retention, pollen shedding ability, silking ability and corn borer tolerance were the criteria used to determine the rows from which ears were selected during the development of LH381.

LH277 and LH185, the progenitors of LH391, are proprietary field corn inbred lines of Holden's Foundation Seeds, L.L.C..

Summer 1998	The inbred line LH277 (a proprietary Holden's inbred) was crossed to the inbred line LH185 (a proprietary Holden's inbred) in Iowa Field/Row 39101-39103.
Winter 1998-99	The S0 seed was grown and self-pollinated in nursery row 24273 in Hawaii.
Summer 1999	S1 ears were grown ear-to-row and self-pollinated in nursery range/row 4476 in Iowa.
Summer 2000	S2 ears were grown ear-to-row and self-pollinated in nursery row 8008 at lowa.
Winter 2000-01	S3 ears were grown ear-to-row and self-pollinated in Hawaii in nursery row 20339.
Summer 2001	S4 ears were grown ear-to-row and self-pollinated in nursery row 53820 at lowa.
Winter 2001-02	S5 ears were grown ear-to-row and self-pollinated in nursery row 11074 at Mexico.
Summer 2002	S6 ears were grown ear-to-row and self-pollinated in nursery row 23867 at lowa.
Summer 2003	S7 ears were grown ear-to-row and self-pollinated in nursery row 28580 at lowa.
Summer 2004	S8 ears were grown ear-to-row and self-pollinated in lowa nursery row 74519-74528.
Winter 2004-05	S9 ears were grown ear-to-row and self-pollinated and final selection made in Hawaii nursery row/field #05KA4A26. Line coded as LH391.

EXHIBIT A (cont'd)

Statement of Stability and Uniformity

LH391 has shown uniformity and stability for all traits described in Exhibit C. It has been self-pollinated and ear-rowed for four generations, with careful attention to uniformity of plant type to ensure homozygosity and phenotypic stability.

Statement of Variants

The line is stable, uniform and no variant traits have been observed or are anticipated in LH391.

EXHIBIT B

Statement of Distinctness

Holden's Foundation Seeds L.L.C. believes that Corn Variety LH391 is most similar to Corn Variety LH277, an inbred developed by Holden's Foundation Seeds L.L.C.

Corn Variety LH391 differ from Corn Variety LH277 at the following traits:

Trait	LH391	LH277
Anther Color	Green-Yellow	Purple
Cob Color	(2.5 GY 8/8) White	(5 RP 5/6) Red
	(Lighter than 2.5 Y 9/2)	(2.5 R 4/8)
Ear Position	Pendent	Upright

Corn variety LH391 has green-yellow anther color, a white cob and pendent ear while comparative corn variety LH277 has purple anther color, a red cob and an upright ear position.

Description of Experimental Design

The corn varieties LH391, LH277 and CM105 were grown at the Waterman, IL observation nursery in years 2005-2006. The varieties were planted in 2 row plots with 15 plants per row in each of the three years. Trait data were collected on 10 random representative plants for most traits from each 2 row plot. Data on qualitative traits are usually collected on 10 plants from each 2 row plot. For Exhibit C all data were pooled and reported as means across the years for subject variety and the standard variety with standard deviation. The varieties are randomly planted in a 4.5 acre observation nursery which is located within a larger 18 acre field. Besides the observation nursery, this field consists of a research seed increase nursery and an IP seed inventory nursery. The location of each of these individual nurseries is rotated each year to a different location within the 18 acre field. Therefore subject inbreds are not planted adjacent to comparative or standard varieties and may be located in different areas of the larger field each year, therefore being influenced by spacial differences within the field. Growing conditions within the field are not uniform as there are some slight topographical variations such as lower areas which may accumulate and retain water or higher areas which are usually drier. The field is tiled and therefore a variety maybe planted close to a tile line while a comparative variety maybe planted further away and in a low spot within the field. Temporal varieties can exist as weather conditions from year to year can vary as well as planting dates can vary from year to year based on weather conditions. Weather conditions each year can vary the maturity rate of the varieties due to either favorable or unfavorable growing conditions.

Trait variability is not observed for each variety within its own test plot-plants are usually uniform and data are collected on the "most" representative plants- variability occurs due to spacial location of the test plot for that variety from year to year and to the temporal variation of weather conditions from year to year during the 2-3 years data are collected.

EXHIBIT B (cont'd)

Waterman Research Station Weather Data 2005-2006

Date	Average Precip. (mm)	Ave. Monthly Temp – Max. (F°)	Ave. Monthly Temp-Min (F°)	Ave. Monthly Rel. Humid Max (%)	Ave. Monthly Rel. Humid – Min (%)
June 2005	0.9	84.7	61.3	89.8	41.7
July 2005	2.0	84.9	61.7	93.4	44.7
August 2005	2.5	82.6	60.4	94.9	50.0
Sept 2005	1.8	79.9	55.0	94.3	44.3
June 2006	2.7	78.4	56.7	89.8	45.9
July 2006	2.3	84.2	64.6	93.5	55.4
August 2006	2.1	87.2	67.5	94.7	57.1
Sept 2006	1.6	80.0	61.6	90.1	50.8

United States Department of Agriculture, Agricultural Marketing Service Science and Technology, Plant Variety Protection Office National Agricultural Library Building, Room 400 Beltsville, MD 20705-2351

OBJECTIVE DESCRIPTION OF VARIETY CORN (Zea mays L.)

., .,			Variety Seed Source	Variety	Name or Temporary Des	ianation
Holden's Foundation Seed L.L.C.			Vallety	LH391	ignation	
77.17	No., City, State, Zip Code and Cou	intrv)	,	EOR OF		PO Number
8350 Minnegan Road, Waterman		may,			I	
				á	20060003	36
Place the appropriate number than necessary. Completeness should	at describes the varietal characters d be striven for to establish an adec	typical of this inbred variety quate variety description.	in the spaces below. Rig	ght justify whole nu	mbers by adding leading	zeroes if
COLOR CHOICES (Use in conjur	nction with Munsell color code to de	escribe all color choices; des	cribe #25 and #26 in Co	mments section):		
01=Light Green 02=Medium Green	06≃Pale Yellow 07≃Yellow	11=Pink 12=Light Red	16=Pale 17=Purpl		21=Buff 22=Tan	
03=Dark Green	08=Yellow-Orange	13=Cherry Red	18=Color		23=Brown	
04=Very Dark Green 05=Green-Yellow	09≃Salmon 10≕Pink-Orange	14=Red	19=White		24=Bronze	
55 5753177611014	ro-Fink-Otalige	15=Red & White	20=White	Capped	25=Variegated (Des 26=Other (Describ	icribe) pe)
STANDARD INBRED CHOICES	(Use the most similar (in backgrou	und and maturity) of these to	make comparisons base	ed on grow-out tri	ai data):	····
Yellow Dent Families: Family Members		Yellow Dent (Unrelated Co109, ND246,	d):	Sweet (Corn: 13, lowa5125, P39, 213:	2
B14 CM105, A63		Oh7, T232		0	10, 104420120, 1 00, 210.	_
B37 B37, B76, H B73 N192, A679.	84 B73, NC268	W117, W153R W182BN		Popcori	n: G1533, 4722, HP301, HI	D7044
C103 Mo17, Va10	2, Va35, A682	44 (OZDI4		3	G 1995, 4722, HF301, H	F/211
Oh43 A619, MS71 WF9 W64A, A554	, H99, Va26 I, A654, Pa91	White Dent: Cl66, H105, Kv2	28	Pipecor	n: o15W, Mo16W, Mo24W	ı
I. TYPE: (describe intermediate t	types in Comments section)	0.55, 11105, 132		Standard Inbred		
2 1=Sweet 2=Dent 3=Flir	nt 4=Flour 5=Pop 6=Ornamental	7=Pipecorn		2 Type		
			Standard Seed Source			
2 1=Northwest 2=North	central 3=Northeast 4=Southea	set 5-South control 6-Se	th		ouice	
	The state of the s	ist 5-30dili Cellilai 0-30	numwest /=Other	2 Region		
	······································	·	outhwest 7=Other	2 Region		
	laptability; show Heat Unit formula	·	outniwest 7=Other		HEAT UNITS	
i. MATURITY (In Region Best Ad DAYS HEAT UNITS	laptability; show Heat Unit formula	in "Comments" section):	nuinwest /=Other	DAYS 68	HEAT UNITS 1400.5	
B. MATURITY (In Region Best Ad DAYS HEAT UNITS 78 1 5 8 8. 0	laptability; show Heat Unit formula	in "Comments" section): ts in silk	nuinwest /=Other	DAYS		
3. MATURITY (In Region Best Ad DAYS HEAT UNITS 78 1588.0 74 1401.5	laptability; show Heat Unit formula 3 From emergence to 50% of plan	in "Comments" section): ts in silk	Juliwest 7-Other	DAYS 68	1400.5	
B. MATURITY (In Region Best Ad DAYS HEAT UNITS 78 1588.0	laptability; show Heat Unit formula S From emergence to 50% of plan From emergence to 50% of plan	in "Comments" section): ts in silk ts in pollen	nuinwest /=Other	DAYS 68	1400.5	
B. MATURITY (In Region Best Ad DAYS HEAT UNITS 78 1588.0 74 1401.5	laptability; show Heat Unit formula S From emergence to 50% of plan From emergence to 50% of plan From 10% to 90% pollen shed	in "Comments" section): ts in silk ts in pollen e quality	Juliwest 7-Other	DAYS 68	1400.5	
B. MATURITY (In Region Best Ad DAYS HEAT UNITS 78 1588.0 74 1401.5	laptability; show Heat Unit formula From emergence to 50% of plan From emergence to 50% of plan From 10% to 90% pollen shed From 50% silk to optimum edible	in "Comments" section): ts in silk ts in pollen e quality	Sample Size	DAYS 68	1400.5	Sample Size
B. MATURITY (In Region Best Ad DAYS HEAT UNITS 78 1588.0 74 1401.5	laptability; show Heat Unit formula From emergence to 50% of plan From emergence to 50% of plan From 10% to 90% pollen shed From 50% silk to optimum edible From 50% silk to harvest at 25%	in "Comments" section): ts in silk ts in pollen e quality 6 moisture		DAYS 68 64 ———	1400.5 1292.5 ————————————————————————————————————	Sample Size
B. MATURITY (In Region Best Ad DAYS HEAT UNITS 78 1588.0 74 1401.5	laptability; show Heat Unit formula From emergence to 50% of plan From emergence to 50% of plan From 10% to 90% pollen shed From 50% silk to optimum edible From 50% silk to harvest at 25%	in "Comments" section): ts in silk ts in pollen e quality 6 moisture Standard Deviation	Sample Size	DAYS 68 64 ———— ————	1400.5 1292.5 Standard Deviation	·
B. MATURITY (In Region Best Ad DAYS HEAT UNITS 78 1588.0 74 1401.5	laptability; show Heat Unit formula From emergence to 50% of plan From emergence to 50% of plan From 10% to 90% pollen shed From 50% silk to optimum edible From 50% silk to harvest at 25% (to tassel tip) Debase of top ear node)	in "Comments" section): ts in silk ts in pollen e quality 6 moisture Standard Deviation 11.8	Sample Size 30	DAYS 68 64 ———— ———— Mean 160.5	1400.5 1292.5 Standard Deviation 24.6	30
B. MATURITY (In Region Best Ad DAYS HEAT UNITS 78 1588.0 74 1401.5	laptability; show Heat Unit formula From emergence to 50% of plan From 10% to 90% pollen shed From 50% silk to optimum edible From 50% silk to harvest at 25% (to tassel tip) to base of top ear node)	in "Comments" section): ts in silk ts in pollen e quality 6 moisture Standard Deviation 11.8 8.8	Sample Size 30 30	DAYS 68 64 Mean 160.5	1400.5 1292.5 Standard Deviation 24.6 12.0	30 30
B. MATURITY (In Region Best Ad DAYS HEAT UNITS 78 1588.0 74 1401.5	laptability; show Heat Unit formula From emergence to 50% of plan From emergence to 50% of plan From 10% to 90% pollen shed From 50% silk to optimum edible From 50% silk to harvest at 25% (to tassel tip) Debase of top ear node) Deformed to the same of the plan to the same of the same of the plan to the same of the plan to the same of the plan to the same of the s	in "Comments" section): ts in silk ts in pollen e quality 6 moisture Standard Deviation 11.8 8.8	Sample Size 30 30	DAYS 68 64 Mean 160.5	1400.5 1292.5 Standard Deviation 24.6 12.0	30 30
DAYS HEAT UNITS 7 8 1 5 8 8. 0 7 4 1 4 0 1. 5 PLANT: 1 9 9. 4 cm Plant Height (to 1 6. 7 cm Length of Top Average Number	laptability; show Heat Unit formula From emergence to 50% of plan From emergence to 50% of plan From 10% to 90% pollen shed From 50% silk to optimum edible From 50% silk to harvest at 25% (to tassel tip) Debase of top ear node) Deformed to the same of the plan to the same of the same of the plan to the same of the plan to the same of the plan to the same of the s	in "Comments" section): ts in silk ts in pollen e quality 6 moisture Standard Deviation 11.8 8.8 2.7 0.0	Sample Size 30 30 30 ———	DAYS 68 64 Mean 160.5 49.4 11.7	1400.5 1292.5	30 30 30

200600036

Application Variety Data		Page 2		Standard Inbred Data		
LEAF:		Standard Deviation	Sample Size	Mean	Standard Deviation	Sample Siz
8.3	cm Width of Ear Node Leaf	0.9	30	7.1	0.7	30
6 7. 1	cm Length of Ear Node Leaf	6.9	30	6 6. 7	9.6	30
5.1	Number of leaves above top ear	0.3	30	5. 7	0.6	15
2 6. 4	degrees Leaf Angle (measure from 2nd leaf above ear at anthes	5.9 is to stalk above leaf)	30	4 7.8	7.0	30
02	Leaf Color (Munsell code 5 GY 4/8)			0 2 (Munsell code 5 GY 4/8)		
1	Leaf Sheath Pubescence (Rate on scale fro	m 1=none to 9=like peach fuzz)		2		
5	Marginal Waves (Rate on scale from 1=non	e to 9=many)		6		
5	Longitudinal Creases (Rate on scale from 1	=none to 9=many)		5		
TASSEL:	W	Standard Deviation	Sample Size	Mean	Standard Deviation	Sample Siz
2.1	Number of Primary Lateral Branches	0.8	30	5. 3	1.2	30
1 6. 5	Branch Angle from Central Spike	8.5	30	3 3.2	9.0	30
3 9. 7	cm Tassel Length (from top leaf collar to tassel tip)	3.1	30	3 4.4	2.6	30
7.4	Pollen Shed (Rate on scale from 0=male ster	ile to 9=heavy shed)		6.2		
	Anther Color (Munsell code 2.5 GY 8/8)	no to similarly alledy		0 7 (Munsell code 2.5 Y 8/10)		
0 2 Glume Color (Munsell code 5 GY 4/8)			1 2 (Munsell code 2.5 R 5/8)			
1		sent		1		
EAR (Unhus	sked Data):				· · · · · · · · · · · · · · · · · · ·	
0.7 Sil	k Color (3 days after emergence) (Munsell cod	e 2.5 Y 8/10)		0 7 (Muns	ell code 2.5 Y 8/10)	
0 2 Fresh Husk Color (25 days after 50% silking) (Munsell code 5 GY 4/8)			0 2 (Munsell code 5 GY 4/8)			
	Husk Color (65 days after 50% Silking) (Muns			2 1 (Munsell code 2.5 Y 8/4)		
	sition of Ear at Dry Husk Stage: 1=Upright 2=H			1	· · · · · · · · · · · · · · · · · · ·	
	sk Tightness (Rate on scale from 1=very loose		•	9		
1 Hus tip) 4=Ver	sk Extension (at harvest): 1=Short (ears expose y Long (>10 cm)	ed) 2=Medium (<8 cm) 3=Long (8-	10 cm beyond ear	1		
EAR (Huske	d Ear Data):	Standard Deviation	Sample Size	Mean	Standard Deviation	Sample Size
1 4, 5	cm Ear Length	0.8	30	1 4.0	1.6	30
3 7. 8	mm Ear Diameter at mid-point	1.2	30	3 8. 0	1.4	15
1 0 6.3	gm Ear Weight	3.6	30	7 3.2	1.9	15
12.7	Number of Kernel Rows	1.0	30	1 3.9	0.9	15
2	Kernel Rows: 1=Indistinct 2=Distinct			2		
1	Row Alignment: 1=Straight 2=Slightly Curved	3=Spiral		1		
9.6	cm Shank Length	2.3	30	6. 8	2.0	15
2	Ear Taper: 1=Slight 2=Average 3=Extreme			2		
	ication Variety Data					

	Page 3		Standard Inbred Da	ala	
8. KERNEL (Dried): S	tandard Deviation	Sample Size	Mean	Standard Deviation	Sample Size
1 0 .5 mm Kernel Length 0.	.5	30	0 9.2	1.0	15
8.5 mm Kernel Width 0.	.7	30	0 8.2	0.6	15
4 . 6 mm Kernel Thickness 0.	.9	30	0 5.1	1.2	15
42.1 % Round Kernels (Shape Grade) 2.	.6	500g	5 6.8	2.6	500g
Aleurone Color Pattern: 1=Homozygous 2=Segregating ((describe)		1		
1 9 Aleurone Color (Munsell code Lighter than 2.5 Y 9/2)			1 9 (Munsell	code Lighter Than 2.5	Y 9/2)
0 7 Hard Endosperm Color (Munsell code 5 Y 8/12)			0.7 (Munsell o	code 2.5 Y 8/8)	
3 Endosperm Type: 1=Sweet (su1) 2=Extra Sweet (sh2) 5=Waxy Starch 6=High Protein 7=High Lysine 10=Other	3=Normal Starch 4=Hig 8=Super Sweet (se)	h Amylose Starch 9=High Oil	0 3		
2 0.4 gm Weight per 100 Kernels (unsized sample) 1.	.1	2200 seeds	2 2.5	1.3	2000 seeds
9. COB: Si	tandard Deviation	Sample Size	Mean	Standard Deviation	Sample Size
1 9 .4 mm Cob Diameter at mid-point 1.	2	30	2 6. 2	1.3	15
1 9 Cob Color (Munsell code Lighter than 2.5 Y 9/2)			1 4 (Munsell	code 5 R 4/10)	
DISEASE RESISTANCE (Rate from 1 (most susceptible) to 9 (most resistance or Strain Options blank if polygenic): A. Leaf Blights, Wilts, and Local Infection Diseases Anthracnose Leaf Blight (Colletotrichum graminicola)	istant); leave blank if not t	ested; leave	7 Anthracnose Le	eaf Blight	
Common Rust (Puccinia sorghi) Common Smut (Ustilago maydis) Eyespot (Kabatiella zeae) Goss's Wilt (Clavibacter michiganense spp. nebraskense) Gray Leaf Spot (Cercospora zeae-maydis) Helminthosporium Leaf Spot (Bipolaris zeicola)	ace _ ace _ ace _		Common Rust Common Smul Eyespot Goss's Wilt Gray Leaf Spot Helminthosporiu Northern Leaf B Southern Leaf Southern Rust Stewart's Wilt	· ·	Race 1 Race O
B. Systemic Diseases Corn Lethal Necrosis (MCMV and MDMV) Head Smut (Sphacelotheca reiliana) Maize Chlorotic Dwarf Virus (MCDV) Maize Chlorotic Mottle Virus (MCMV) Maize Dwarf Mosaic Virus (MDMV)	rain			c Dwarf Virus Mottle Virus Iosaic Virus In Mildew of Corn	Strain
Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Fusarium moniliforme) Gibberella Stalk Rot (Gibberella zeae) Other (Specify) D. Ear and Kernel Rots			Anthracnose S Diplodia Stalk F Fusarium Stalk Gibberella Stal Other (Specify)	Rot : Rot k Rot	
Aspergillus Ear and Kernel Rot (<i>Aspergillus flavus</i>) Diplodia Ear Rot (<i>Stenocarpella maydis</i>) Fusarium Ear and Kernel Rot (<i>Fusarium moniliforme</i>) Gibberella Ear Rot (<i>Gibberella zeae</i>) Other (Specify)			Aspergillus Ear Diplodia Ear Ro Fusarium Ear 8 Gibberella Ear Other (Specify)	ot & Kernel Rot Rot	
Application Variety Data Note: Use chart on first page to choose color codes for color traits.			Standard Inbred Da	ıta	

200600036

Application Variety Data	Page 4		Standard Inbred Data		
11. INSECT RESISTANCE (Rate from 1 (most susceptible) to 9 (most resistant);					
leave blank if not tested): Standard D	Deviation	Sample Size	Standard Deviation Sample Size		
Banks Grass Mite (Oligonychus pratensis)			Banks Grass Mite		
Corn Earworm (<i>Helicoverpa zea</i>) Leaf-Feeding			Corn Earworm Leaf Feeding		
Silk Feeding : mg larval wt. Ear Damage	<u> </u>		Ear Damage		
Corn Leaf Aphid (Rhopalosiphum maidis)			Corn Leaf Aphid		
Corn Sap Beetle (Carpophilus dimidiatus)	·		Corn Sap Beetle		
European Corn Borer (Ostrinia nubilalis) 1st Generation (Typically Whorl Leaf Feeding)			European Corn Borer		
2nd Generation (Typically Leaf Sheath-Collar Feeding) Stalk Tunneling: cm tunneled/plant			1st Generation 2nd Generation		
Fall Armyworm (Spodoptera frugiperda)	_ ·				
Leaf-Feeding			Fall Armyworm Leaf Feeding		
	 -				
Maize Weevil (Sitophilus zeamaize) Northern Rootworm (Diabrotica barberi)			Maize Weevil Northern Rootworm		
Southern Rootworm (Diabrotica undecimpunctata)			Southern Rootworm		
Southwestern Corn Borer (<i>Diatraea grandiosella</i>) Leaf Feeding			Southwestern Corn Borer Leaf Feeding		
Stalk Tunneling : cm tunneled/plant	- -				
Two-spotted Spider Mite (Tetranychus urticae) Western Rootworm (Diabrotica virgifera virgifera)			Two-spotted Spider Mite Western Rootworm		
Other (Specify)			Other (Specify)		
12. AGRONOMIC TRAITS:			*****		
7 Stay Green (at 65 days after anthesis) (Rate on a scale from 1=worst to 9	9=excellent.)		1 Stay Green		
0 0 . 0 % Dropped Ears (at 65 days after anthesis)	•		0 0 . 1 % Dropped ears		
0 0 .0 % Pre-anthesis Brittle Snapping			0 0 . 0 % Pre-anthesis Brittle Snapping		
0 0. 0 % Pre-anthesis Root Lodging			0 0 . 0 % Pre-anthesis Root Lodging		
0 0. 0 % Post-anthesis Root Lodging (at 65 days after anthesis)			0 0 . 0 % Post-anthesis Root Lodging		
Kg/ha Yield of Inbred Per Se (at 12-13% grain moisture)		Ī	Yield		
13. MOLECULAR MARKERS: (0=data unavailable; 1=data available but not supplied;	; 2=data supplied)				
0 Isozymes 0 RFLP's 0 RAPD'sOth	ner (Specify)				
REFERENCES:					
Butler, D.R. 1954. A System for the Classification of Corn Inbred Lines. PhD Thesis,	Ohio State Univers	situ			
Emerson, R.A., G.W. Beadle, and A.C. Fraser. 1935. A Summary of Linkage Studies	in Maize, Cornell	A.E.S., Mem. 180.	District Control of the Control of t		
Farr, D.F., G.F. Bills, G.P. Chamuris, A.Y. Rossman. 1989. Fungi on Plant and Plant Inglett, G.E. (Ed.) 1970. Corn: Culture, Processing, Products. Avi Publishing Compan	v. Westport, CT.		merican Phytopathological Society, St. Paul, MN.		
Jugenheimer, R.W. 1976. Corn: Improvement, Seed Production, and Uses. John Wilk McGee, D.C. 1988. Maize Diseases. APS Press, St. Paul, MN. 150 pp.	ey & Sons, New Yo	ork.			
Munsell Color Chart for Plant Tissues. Macbeth. P.O. Box 230. Newburgh, N.Y. 1255	1-0230				
The Mutants of Maize. 1968. Crop Science Society of America. Madison, WI. Shurtleff, M.C. 1980. Compendium of Com Diseases. APS Press, St. Paul, MN. 105					
Sprague, G.F., and J.W. Dudley (Editors). 1988. Corn and Corn Improvement, Third I	Edition. Agronomy	Monograph 18. As	SA, CSSA, SSSA, Madison, WI.		
Stringfield, G.H. Maize Inbred Lines of Ohio. Ohio A.E.S., Bul. 831, 1959. U.S. Department of Agriculture. 1936, 1937. Yearbook.					
COMMENTS (e.g. state how heat units were calculated, standard inbred seed source,	, and/or where data	a was collected. C	ontinue in Exhibit D):		
Heat Unit Calculation: GDU = Daily Max Temp (<=86°F) + D	Daily Min Ten	np (>=50°F) -	- 50°F		
	Heat Unit Calculation: GDU = <u>Daily Max Temp (<=86°F) + Daily Min Temp (>=50°F)</u> - 50°F				
Supplemental data provided for pollen shed, ear weight, % round kemels a inventory data. Supplemental data for quantitative traits for the subject vari	na weight per 10 iety 'LH391' prov	ided from seed	inventory 2006 and 2007 and production parent test		
data.	•		•		

REPRODUCE LOCALLY Include form quarte		all reproductions.	FORM APPROVED - OMS No. 0581/00
U.S. DEPARTMENT OF AGRICULTURAL MARKETING EXHIBIT E	SERVICE	Application is required in order to certificate is to be issued (7 U.S.) confidential until the certificate is	determine if a plant variety protection 2. 2421). The information is hidd
STATEMENT OF THE BASIS OF 1. NAME OF APPLICANT(S)	OWNERSHIP	A TOWNS AND DESIGNATION	
		Z TEMPORARY BESIGNATION OR EXPERIMENTAL NUMBER	
Holden's Foundation Seeds L	L.C.	and a second processing the second processin	LH391
4. ADDRESS (street out No., cr.8. F.O. No., City. Stells	. and III. don't Country	5. TELEPHONE (Indice 3700 exce)	S. FAX Oneluses area code;
8350 Minnegan Road		(815) 758-9281	(815) 758-3117
Waterman, IL 60556 U.S.A.		7. PVPO NUMBER	200500036
8. Does the applicant own all rights to the vari 9. Is the applicant (individual or company) a U			
			<u> </u>
10. Is the applicant the original owner?	YES NO	If no, please answer <u>one</u> of the	following:
a. If the original rights to variety were own	ed by individual(s), is	(are) the original owner(s) a U.S. Nati	ional(s)?
	YES NO	If no, give name of country	
b. If the original rights to variety were own	ned by a company(les)), is (are) the original owner(s) a U.S.	based company?
	YES NO	If no, give name of country	,
11. Additional explanation on ownership (if nee	eded, use the reverse	for extra space):	
Foundation Seeds, L.L.C. breeder, all rights to any in	By agreement be nvention, discover	veloped by a breeder employe etween Holden's Foundation S ry or development are assigne n invention, discovery or devel	Seeds, L.L.C. and the d to Holden's
PLEASE NOTE:	· ·		
Plant variety protection can only be afforded to	the owners (not licen	sees) who meet the following criteria:	•
If the rights to the variety are owned by the o national of a country which affords similar pro	original breeder, that o	erson must be a U.S. national, nation	al of a UPOV member country, or
If the rights to the variety are owned by the c nationals of a UPOV member country, or own genus and species.	ompany which emplo	ved the original breeder(s), the compa	any must be U.S. based, owned by
3. If the applicant is an owner who is not the ori	ginal owner, both the	original owner and the applicant mus	t meet one of the above criteria.
The original breeder/owner may be the individu Act for definitions.			
According to the Paperwork Reduction Act of 1995, an agency control number. The valid OMB control number for this informates response, including the time for reviewing the instructions, sear	ition collection is 0581-0055.	The lime required to complete this information or	illection is estimated to average 6 minutes per

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, or marital or family status. (Not all prohibited bases apply to all programs). Persons with disabilities who require alternative means for communication of program information (braille, large print, audiolape, stc.) should contact the USDA's TARGET Center at 202-720-2600 (voice and TDD). To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5984 (voice and TDD). USDA is an equal opportunity provider and employer.

ST-470-E (04-99) (Destroy previous editions).

REPRODUCE LOCALLY, Include form number and date on all reproductions.

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 5 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY **PLANT VARIETY PROTECTION OFFICE** BELTSVILLE, MD 20705

EXHIBIT F

NAME OF OWNER (S)	ADDRESS (Street and No. or RD No., City, State, and Zip Code and Country)	TEMPORARY OR EXPERIMENTAL DESIGNATION
Holden's Foundation Seeds LLC	8350 Minnegan Road, Waterman, IL 60556 USA	
		VARIETY NAME LH391
NAME OF OWNER REPRESENTATIVE (S)	ADDRESS (Street and No. or RD No., City, State, and Zip Code and Country)	Echecopera, islamic property and an analysis of the control of the
Timothy R. Kain	8350 Minnegan Road, Waterman, IL 60556 USA	PVPO NUMBER
		200600036

I do hereby declare that during the life of the certificate a viable sample of propagating material of the subject variety will be deposited, and replenished as needed periodically, in a public repository in the United States in accordance with the regulations established by the Plant Variety Protection Office.

moth R. Le